

PFA-100[®] Analyzer Bibliography

von Willebrand Disease Assessment

Fressinaud E, Veyradier A, Truchaud F, Martin I, Boyer-Neumann C, Trossaert M, Meyer D: **Screening for von Willebrand Disease with a new analyzer using high shear stress: A study of 60 cases.** Blood 1998; 91:1325-1331.

Di Paola J; Federici AB; Mannucci PM; Canciani MT; Kritzik M; Kunicki TJ; Nugent D: **Low platelet alpha sub(2) beta sub(1) levels in type I von Willebrand disease correlate with impaired platelet function in a high shear stress system.** Blood 1999; 93(11):3578-3582.

Cattaneo M, Federici AB, Lecchi A, Agati B, Lombardi R, Stabile F, Bucciarelli P. **Evaluation of the PFA-100[®] system in the diagnosis and therapeutic monitoring of patients with von Willebrand disease.** Thromb Haemost 1999;82(1):35-39.

Fressinaud E, Veyradier A, Sigaud M, Boyer-Neumann C, Le Boterff C, Meyer D. **Therapeutic monitoring of von Willebrand disease: Interest and limits of a platelet function analyser at high shear rates.** British Journal of Haematology 1999;106(3):777-783.

Favoloro EJ, Facey D, Henniker A. **Use of a novel platelet function analyzer (PFA-100[™]) with high sensitivity to disturbances in von Willebrand Factor to screen for von Willebrand's Disease and other disorders.** Am. J Hematol 1999;62:165-174.

Sadler J, Manucci P, Berntrop N, Bochkov N, Boulyjenkov V, Ginsburg D, Meyer D, Peake I, Rodeghiero F, Siravastava A: **Impact, diagnosis and treatment of von Willebrand disease.** Thromb Haemost 2000; 84: 160-174.

Veyradier A, Fressinaud E, Boyer-Neumann C, Trossaert M, Meyer D: **von Willebrand factor ristocetin cofactor activity correlates with the platelet function in a high shear stress system,** Thromb Haemost 2000, 84: 27-728.

Schlamadinger A, Kerenyi A, Muszbek L, Boda Z: **Comparison of the O'Brian filter test and the PFA-100 platelet analyzer in the laboratory diagnosis of von Willebrand's disease.** Thromb Haemost 2000; 84(1): 88-92

Favoloro E, Kershaw G, Bukuya M, Hertzberg M, Koutts J: **Laboratory diagnosis of von Willebrand disorder (vWD) and monitoring of DDAVP therapy: efficacy of the PFA-100 and vWF:CBA as combined diagnostic strategies;** Haemophilia 2001; 7: 180-189.

Frank RD, Kunz D, Wirtz DC: **Acquired von Willebrand disease-hemostatic management of major orthopedic surgery with high-dose immunoglobulin, desmopressin and continuous factor concentrate infusion:** Am J Hematol 2002; 70:64-71.

Franchini M, Gandini G, Manzato F, Lippi G: **Evaluation of the PFA-100 system for monitoring desmopressin therapy in patients with type 1 von Willebrand disease;** Haematologica 2002; 6:270.

De Jonge E, Levi M, Bueller HR, Berends F, Kesecioglu J: **Decreased circulating levels of von Willebrand factor after intravenous administration of a rapidly degradable hydroxyethyl starch (HES 200/0.5/6) in healthy human subjects;** Intensive Care Med 2001; 27:1825-1829.